

Supplementary Table 2. Outcomes by treatment modality: subgroup analysis of posterior circulation, NIHSS ≥6

Characteristic	SMT (n=12, 22%)	MT (n=42, 78%)	P
Postprocedural any ICH	5 (42)	12 (29)	0.49
Post procedural sICH	2 (17)	4 (9.5)	0.60
Classification of ICH			0.39
None	9 (75)	37 (88)	
HI1	1 (8.3)	2 (4.8)	
HI2	1 (8.3)	1 (2.4)	
PH2	1 (8.3)	1 (2.4)	
SAH	0 (0)	1 (2.4)	
Cerebral edema in same arterial territory with midline shift	3 (25)	9 (21)	>0.99
Suboccipital-hemicraniotomy	1 (8.3)	2 (4.8)	0.54
Predischarge NIHSS	27.5±12.3 (0–42)	18±15.8 (0–42)	0.07
Median (IQR)	29 (22–36)	12 (5–33)	
Predischarge mRS	3.5±2 (0–6)	4±1.4 (0–6)	0.03
Median (IQR)	6 (4.4–6)	4.5 (4–6)	
Length of stay (day)	7.8±7.1 (1–26)	13.4±14.1 (1–60)	0.19
Median (IQR)	5 (3–12)	8 (4–17)	
Discharge destination			0.09
Home	1 (8.3)	3 (7.1)	
Acute rehabilitation/nursing home	2 (17)	22 (52)	
Long-term acute care hospitals	0 (0)	1 (2.4)	
Death	9 (75)	16 (38)	
90-day mRS	5.4±1.4 (2–6)	4±1.9 (1–6)	0.03
mRS 0–2 at 90 days	1 (9.1)	10 (27)	0.42
Mortality at 90 days	9 (82)	15 (41)	0.02

Data are presented as n (%) or mean±standard deviation (range) unless otherwise indicated. Person chi-squared test, Fisher's exact test compare categorical variables; Wilcoxon rank-sum test for continuous variables; false discovery rate correction for multiple testing.

NIHSS, National Institutes of Health Stroke Scale; SMT, standard medical therapy; MT, mechanical thrombectomy; ICH, intracranial hemorrhage; sICH, symptomatic intracranial hemorrhage; HI1, hemorrhagic infarction type 1; HI2, hemorrhagic infarction type 2; PH1, parenchymal hematoma type 1; PH2, parenchymal hematoma type 2; SAH, subarachnoid hemorrhage; IQR, interquartile range; mRS, modified Rankin Scale.