

Supplementary Table 2. Periprocedural details of the included patients before matching

Variable	Overall (n=1,057)	MT-IVT (n=715)	IVT alone (n=342)	P
First line technique				
Aspiration	-	147 (21)	NA	
Both	-	418 (61)	NA	
Stent retriever	-	122 (18)	NA	
Side				0.14
Right	486 (48)	349 (50)	137 (44)	
Left	526 (52)	354 (50)	172 (56)	
Mothership versus drip and ship				
Drip and ship	341 (37)	299 (43)	42 (19)	<0.001***
Mothership	582 (63)	399 (57)	183 (81)	
Onset to IVT needle time (min)	150 (104–225)	145 (100–210)	164 (115–247)	0.007**
Onset to arterial puncture (min)	-	225 (161–322)	NA	
Puncture to recanalization time (min)	-	37 (25–58)	NA	
Onset to recanalization (min)	-	279 (204–376)	NA	
Anesthesia				
CS/LA	-	487 (69)	NA	
GA	-	217 (31)	NA	
Puncture site				
Femoral	-	442 (95)	NA	0.47
Radial	-	21 (4.5)	NA	
Imaging after MT				
CT	-	463 (67)	NA	0.16
Both	-	119 (17)	NA	
MRI	-	108 (16)	NA	
No imaging	-	1 (0.1)	NA	

Values are presented as n (%) or median (interquartile range).

MT, mechanical thrombectomy; IVT, intravenous thrombolysis; CS/LA, conscious sedation/local anesthesia; GA, general anesthesia; CT, computed tomography; MRI, magnetic resonance imaging; NA, not applicable.

** $P < 0.01$; *** $P < 0.001$.