- GWAS Catalog	2,839.		«<-+>»	TOP
	rs2237892 (Body mass index)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
- GRASP			«<-+>»	TOP
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (HOMA-B)			
	rs2237892 (Body mass index (E	BMI))		
	rs2237892 (Corrected insulin re	esponse (CIR))		
	rs2237892 (Type 2 diabetes)			
	rs2237892 (2 hour glucose)			
	rs2237892 (HOMA-B)			
	rs2237892 (Fasting insulin)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (HOMA-IR)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)			
	rs2237892 (Type 2 diabetes)	FDNAID1 in blood)		
	rs2237892 (Gene expression of			
	rs2237892 (Gene expression of rs2237892 (Type 2 diabetes)	TIFKE III blood)		
	rs2237892 (Type 2 diabetes)	obaca individuals (PMI)	30 Ka(m2))	
	rs2237892 (Type 2 diabetes in t			
	rs2237892 (Type 2 diabetes)		/ (g/mz/)	
	rs2237892 (Coronary artery dis			
	rs2237892 (Colonary artery dis	Care (CRD/)		
	"szeriest (type z diabetes)			

Supplementary Fig. 2. Genome-wide association study (GWAS) results of rs2237892 from Korean Reference Genome Database (KRGDB). Search result from KRGDB. The original data comes from GWAS catalog and genome-wide repository of associations between SNPs and phenotypes (GRASP) database. The results from GWAS catalog had genome-wide significance ($P < 5 \times 10^{-8}$), while those from GRASP had a range of *P* value from nominal significance to genome-wide significance.