

Supplementary Table 1. Detection rate of distribution of wet serum concentration (pg/mL) of individual compounds belonging to PCBs and OCPs (*n*=323)

POP	Detection rate, %	Min	Percentile, pg/mL			Max
			25th	50th	75th	
PCB						
PCB 118	99.4	0.53	23.42	46.44	110.44	1,007.41
PCB 138	99.7	1.07	59.61	123.97	272.70	2,511.23
PCB 153	99.4	1.07	72.71	146.33	331.41	3,263.27
PCB 180	92.0	1.07	34.22	81.00	206.11	1,811.26
PCB 187	75.5	1.07	12.48	41.43	93.50	958.63
OCP						
β-HCH	64.1	1.07	1.07	52.61	118.22	676.88
p,p'-DDE	98.8	1.07	464.00	878.48	1440.22	8,242.63
p,p'-DDT	69.3	1.07	1.07	42.05	65.83	284.67
Trans-nonachlor	78.3	0.43	15.37	35.66	61.69	219.95

PCB, polychlorinated biphenyl; OCP, organochlorine pesticide; POP, persistent organic pollutant; HCH, hexachlorocyclohexane; DDE, dichlorodiphenyldichloroethylene; DDT, dichlorodiphenyltrichloroethane.