

Supplementary Table 3. The risk of cardiovascular death according to the component of metabolic dysfunction-associated fatty liver disease

Variable	Normal	Obesity (BMI \geq 23 kg/m ²)	Type 2 diabetes mellitus	Lean with metabolic dysfunction
All				
Crude HR	1.0 (ref)	1.17 (0.85–1.60)	1.08 (0.42–2.82)	1.58 (0.84–3.00)
Age-adjusted HR	1.0 (ref)	1.17 (0.85–1.61)	0.99 (0.38–2.57)	1.21 (0.64–2.28)
Multivariable-adjusted ^a HR	1.0 (ref)	1.05 (0.76–1.44)	0.99 (0.37–2.61)	1.21 (0.64–2.30)
Men				
Crude HR	1.0 (ref)	0.95 (0.66–1.36)	1.20 (0.41–3.52)	1.20 (0.56–2.57)
Age-adjusted HR	1.0 (ref)	1.01 (0.70–1.44)	1.20 (0.41–3.52)	1.15 (0.54–2.46)
Multivariable-adjusted ^a HR	1.0 (ref)	0.93 (0.65–1.34)	1.28 (0.43–3.80)	1.16 (0.54–2.50)
Women				
Crude HR	1.0 (ref)	2.08 (1.07–4.07)	0.68 (0.08–6.05)	2.13 (0.69–6.60)
Age-adjusted HR	1.0 (ref)	1.26 (0.64–2.48)	0.54 (0.06–4.83)	1.17 (0.37–3.71)
Multivariable-adjusted ^a HR	1.0 (ref)	1.26 (0.64–2.49)	0.44 (0.05–4.00)	1.18 (0.37–3.77)

BMI, body mass index; HR, hazard ratio.

^aAdjusted for age, daily alcohol consumption, regular physical activity, smoking status, total cholesterol, and statin use.