

Supplementary Table 1. Baseline characteristics at event in patients with type 2 diabetes mellitus and established cardiovascular disease

Variable	Value
Age, yr	70.0±11.9
Male sex	1,371 (66.9)
BMI, kg/m ²	24.7±4.1
Smoking	
Non-smoker	1,178 (57.5)
Former smoker	411 (20.0)
Current smoker	426 (20.8)
Comorbidities	
Hypertension	1,522 (74.2)
History of IHD	690 (33.7)
History of stroke	304 (14.8)
History of PAD	89 (4.3)
History of HF	234 (11.4)
Diabetes duration, yr	12.0±10.1
Range of diabetes duration, yr	
<10	674 (32.9)
10–19	481 (23.5)
≥20	452 (22.0)
Unknown	443 (21.6)
SBP, mm Hg	127.3±17.8
DBP, mm Hg	73.2±11.1
Random plasma glucose, mg/dL	172.5±78.2
HbA1c, %	7.5±1.6
Creatinine, mg/dL	1.40±1.53
eGFR, mL/min/1.73 m ²	
≥60	1,311 (64.0)
45–59	304 (14.8)
30–44	189 (9.2)
<30	234 (11.4)
Total cholesterol, mg/dL	150.0±46.7
Triglycerides, mg/dL	146.5±117.5
HDL-C level, mg/dL	42.8±11.8
LDL-C level, mg/dL	86.7±36.1

Values are presented as number (%) or mean ± standard deviation. BMI, body mass index; IHD, ischemic heart disease; PAD, peripheral artery disease; HF, heart failure; SBP, systolic blood pressure; DBP, diastolic blood pressure; HbA1c, glycosylated hemoglobin; eGFR, estimated glomerular filtration rate; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol.