

Supplementary Table 1. Trends in the prevalence and incidence of diabetes-related ESKD-NHIS database (2008 to 2019)

Variable	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Prevalence rate ^a of diabetes-related ESKD												
≥30 yr	982.0	989.7	988.4	1,004.3	1,030.6	1,052.3	1,082.1	1,103.0	1,126.4	1,135.8	1,146.5	1,169.3
30–59 yr	1,094.9	1,081.4	1,075.0	1,082.1	1,096.6	1,116.4	1,150.4	1,164.3	1,162.2	1,153.8	1,161.7	1,169.2
60–79 yr	962.6	983.7	983.8	1,003.1	1,040.2	1,065.6	1,089.9	1,120.5	1,162.6	1,183.2	1,198.6	1,232.4
≥80 yr	418.2	494.8	514.6	544.2	563.9	587.1	661.4	688.1	758.9	815.9	837.3	902.2
Incidence rate ^a of diabetes-related ESKD												
≥30 yr	235.1	224.5	206.9	213.2	213.3	202.6	207.5	216.1	219.8	207.4	200.1	201.5
30–59 yr	241.7	224.2	213.4	214.4	214.0	199.1	205.4	214.9	208.1	198.6	191.1	187.9
60–79 yr	238.4	228.9	208.2	216.2	217.7	210.2	210.9	219.6	227.9	214.9	207.0	211.1
≥80 yr	170.4	187.7	155.4	185.0	184.3	180.8	209.6	221.2	258.1	244.0	244.8	259.0

ESKD, end-stage kidney disease; NHIS, National Health Insurance Service.

^aAge- and sex-standardized rates.