

Supplementary Table 4. Body composition parameters of patients with and without fatty liver

Variable	Fatty liver group	Non-fatty liver group	P value
Number	100	177	
Age, yr	62.66 ± 10.18	59.79 ± 9.45	0.0195
BMI, kg/m ²	25.57 ± 2.52	23.24 ± 2.99	<0.0001
WC, cm	88.37 ± 7.37	81.72 ± 8.08	<0.0001
VAT, cm ²	137.12 ± 45.43	91.25 ± 45.44	<0.0001
VAT_SUV _{max}	0.31 ± 0.07	0.49 ± 0.15	<0.0001
SAT, cm ²	192.57 ± 59.22	156.68 ± 57.29	<0.0001
SAT_SUV _{max}	0.24 ± 0.06	0.35 ± 0.10	<0.0001
CSMA, cm ²	130.69 ± 34.70	122.65 ± 32.34	0.0539
Muscle_HU	27.64 ± 6.97	28.71 ± 6.72	0.2122
Muscle_SUV _{max}	0.77 ± 0.20	0.73 ± 0.15	<0.0001

Values are presented as mean ± standard deviation. One patient whose liver could not be evaluated due to severe beam-hardening artifact was excluded.

BMI, body mass index; WC, waist circumference; VAT, visceral adipose tissue; SUV_{max}, maximum standardized uptake value; SAT, subcutaneous adipose tissue; CSMA, cross-sectional muscle area; HU, Hounsfield unit.