

Supplementary Table 3. Multivariable hazard ratio and stratified risk analysis for T2DM according to parity, excluding 315 women with gestational diabetes mellitus

No. of live births	No. of events/ Total no.	Incidence rate, / 1,000 person-yr (95% CI)	Absolute risk, %	Crude		Model 1	
				HR (95% CI)	P value	HR (95% CI)	P value
0	646/45,567	1.61 (1.48–1.73)	1.42	Ref		Ref	
1	575/32,267	2.01 (1.85–2.18)	1.78	1.25 (1.12–1.40)	<0.001	1.17 (1.04–1.31)	0.007
2	1,524/106,476	1.61 (1.53–1.69)	1.43	1.00 (0.92–1.10)	0.941	0.89 (0.81–0.97)	0.011
3	798/42,761	2.10 (1.96–2.25)	1.87	1.31 (1.18–1.46)	<0.001	1.10 (0.99–1.22)	0.073
4	307/11,177	3.10 (2.76–3.47)	2.75	1.94 (1.69–2.22)	<0.001	1.52 (1.32–1.74)	<0.001
5 or more	160/3,596	5.05 (4.30–5.89)	4.45	3.16 (2.66–3.76)	<0.001	2.25 (1.88–2.68)	<0.001

Model 1: age and race.

T2DM, type 2 diabetes mellitus; CI, confidence interval; HR, hazard ratio.