

Supplemental Table S3. Multivariate Logistic Regression Analyses for the Association between Metabolic Syndrome and Secondhand Smoke Exposure According to Daily Time, Frequency, and Duration in Never Smokers

Variable	Age- and sex-adjusted	Multivariate
SHS exposure	1.19 (1.13–1.25)	1.09 (1.02–1.16)
Daily time of SHS exposure		
< 1 hr	1.09 (0.99–1.19)	0.91 (0.82–1.01)
≥ 1 hr	1.49 (1.35–1.66)	1.18 (1.04–1.34)
<i>P</i> for trend	<0.001	0.190
Frequency of SHS exposure		
< 3 times/wk	1.06 (0.99–1.13)	1.03 (0.95–1.11)
≥ 3 times/wk	1.42 (1.32–1.54)	1.18 (1.08–1.29)
<i>P</i> for trend	<0.001	0.001
Duration of SHS exposure		
< 10 yr	1.12 (1.01–1.24)	1.01 (0.89–1.13)
≥ 10 yr	1.27 (1.17–1.36)	1.05 (0.96–1.15)
<i>P</i> for trend	<0.001	0.295

Values are expressed as odds ratio (95% confidence interval).

Multivariate model was adjusted for age, sex, body mass index, frequency of alcohol drinking, frequency of vigorous exercise, blood urea nitrogen, creatinine, uric acid, total cholesterol, low-density lipoprotein cholesterol, and high-sensitivity C-reactive protein.

SHS, secondhand smoke.