

Supplemental Table S1. The Number of Subjects According to Glucose Tolerance in the AhRL and MIS-ATP Quartile Groups Aged 70 and 75 Years

Quartile	Age at 70 yr			Age at 75 yr		
	NGT (n=710)	IGT (n=99)	DM (n=102)	NGT (n=562)	IGT (n=93)	DM (n=87)
AhRL						
Q1	180 (79.3)	22 (9.7)	25 (11.0)	140 (78.7)	20 (11.2)	18 (10.1)
Q2	182 (79.8)	23 (10.1)	23 (10.1)	142 (78.0)	20 (11.0)	20 (11.0)
Q3	182 (79.8)	23 (10.1)	23 (10.1)	146 (76.9)	20 (10.5)	24 (12.6)
Q4	166 (72.8)	31 (13.6)	31 (13.6)	134 (69.8)	33 (17.2)	25 (13.0)
MIS-ATP						
Q1	167 (73.6)	30 (13.2)	30 (13.2)	140 (73.3)	22 (11.5)	29 (15.2)
Q2	179 (78.5)	28 (12.3)	21 (9.2)	138 (75.8)	28 (15.4)	16 (8.8)
Q3	182 (79.8)	25 (11.0)	21 (9.2)	135 (74.6)	28 (15.5)	18 (9.9)
Q4	182 (79.8)	16 (7.0)	30 (13.2)	149 (79.2)	15 (8.0)	24 (12.8)

Values are expressed as number (%).

AhRL, arylhydrocarbon receptor ligand activity; MIS-ATP, mitochondrial inhibiting substance activity measured by intracellular ATP content; NGT, normal glucose tolerance; IGT, impaired glucose tolerance; DM, diabetes mellitus.