

Supplemental Table S2. Risk for NAFLD (Fatty Liver Index ≥ 60) According to the Quartile of Average Successive Variability of Bodyweight

	Presence of NAFLD	Model 1	Model 2	Model 3	Model 4
Quartile 1	66 (1.8)	1 (reference)	1 (reference)	1 (reference)	1 (reference)
Quartile 2	118 (3.1)	1.77 (1.30–2.40)	1.57 (1.15–2.14)	1.55 (1.12–2.14)	1.42 (1.01–2.00)
Quartile 3	192 (5.0)	2.94 (2.22–3.91)	2.37 (1.77–3.16)	2.38 (1.76–3.22)	1.96 (1.43–2.70)
Quartile 4	375 (9.8)	6.08 (4.66–7.93)	5.08 (3.87–6.68)	5.00 (3.76–6.66)	2.74 (2.01–3.72)

Values are expressed as number (%) or odds ratio (95% confidence interval). Model 1: No adjustment; Model 2: Adjusted for age, sex; Model 3: Adjusted for Model 2+fasting blood glucose, systolic blood pressure, triglyceride, low-density lipoprotein cholesterol, smoking amount, exercise, and alcohol frequency; Model 4: Adjusted for Model 2+glycated hemoglobin, systolic blood pressure, triglyceride, low-density lipoprotein cholesterol, smoking amount, exercise and alcohol frequency, body mass index, homeostasis model assessment of insulin resistance, aspartate transaminase, and alanine aminotransferase.

NAFLD, nonalcoholic fatty liver disease.