

Supplemental Table S1. Baseline Characteristics of High Mean TT3 Group and Low Mean TT3 Group

Characteristic	SLD risk cohort (n=1,216)	Low mean TT3 group (n=609)	High mean TT3 group (n=607)	P value
Age, yr	43.0 (37.0–50.0)	43.0 (37.0–50.0)	43.0 (37.0–49.0)	0.477
Female sex	432 (35.5)	227 (37.3)	205 (33.8)	0.202
BMI, kg/m ²	23.6 (21.6–25.7)	23.4 (21.5–25.5)	23.7 (21.7–25.8)	0.199
Obesity				0.478
Normal (BMI <23 kg/m ²)	509 (41.9)	265 (43.5)	244 (40.2)	
Overweight (BMI 23–25 kg/m ²)	306 (25.2)	151 (24.8)	155 (25.5)	
Obese (BMI ≥25 kg/m ²)	401 (33.0)	193 (31.7)	208 (34.3)	
TSH level, mU/L	1.74 (1.18–2.39)	1.72 (1.18–2.40)	1.74 (1.20–2.38)	0.962
FT4 level, ng/dL	1.32 (1.21–1.43)	1.31 (1.20–1.43)	1.33 (1.21–1.43)	0.251
TT3 level, ng/dL	107.80 (97.23–118.10)	104.00 (94.56–113.65)	111.80 (101.70–121.60)	<0.001
AST level, IU/L	21.0 (17.0–26.0)	21.0 (17.0–26.0)	21.0 (17.0–26.0)	0.398
ALT level, IU/L	21.0 (15.0–29.0)	20.0 (15.0–29.0)	21.0 (15.0–30.0)	0.188
HbA1c level, %	5.1 (5.0–5.2)	5.1 (5.0–5.2)	5.1 (5.0–5.2)	0.197
Triglyceride level, mg/dL	107.0 (76.0–155.0)	102.0 (76.0–156.0)	109.0 (77.0–154.0)	0.775
HDL-C level, mg/dL	51.0 (43.0–61.0)	52.0 (44.0–62.0)	50.0 (43.0–60.0)	0.217
LDL-C level, mg/dL	116.0 (96.0–136.0)	117.0 (98.0–137.0)	115.0 (94.0–135.0)	0.446
Median follow-up, mo	63.0 (51.2–71.0)	63.0 (51.0–71.0)	63.0 (52.0–71.0)	0.847

Values are expressed as median (interquartile range) or number (%).

TT3, total triiodothyronine; SLD, steatotic liver disease; BMI, body mass index; TSH, thyroid stimulating hormone; FT4, free thyroxine; AST, aspartate aminotransferase; ALT, alanine transaminase; HbA1c, hemoglobin A1c; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol.