

Supplemental Table S3. Analysis of PFS Showing the HR and Discrimination of the Original G Compartment, the S-GRAS^a Score, and Groups of S-GRAS Score in Patients with Available Data of Ki67 Index (*n*=79)^b

Scores	Variable	No. (%)	No. of relapse (%)	HR (95% CI)	<i>P</i> value	Harrell's C-index (95% CI) ^c	<i>P</i> value ^c	<i>P</i> value ^d
G	Ki67 index, %	79 (100.0)	46 (58.2)			0.648 (0.552–0.731) ^e	<0.001 ^e	<0.001 ^e
0	0–9	36 (45.6)	17 (27.8)	Ref				
1	10–19	23 (29.1)	14 (60.9)	1.62 (0.80–3.30)	0.180			
2	≥20	20 (25.3)	15 (75.0)	2.99 (1.48–6.10) ^e	0.003 ^e			
S-GRAS		79 (100.0)	46 (58.2)	1.76 (1.50–2.06) ^e	<0.001 ^e	0.805 (0.720–0.870) ^e	Ref	
S-GRAS group		79 (100.0)	46 (58.2)			0.803 (0.717–0.864) ^e		Ref
0–1		23 (29.1)	9 (39.1)	Ref				
2–3		29 (36.7)	13 (44.8)	1.30 (0.55–3.06)	0.546			
4–5		7 (8.9)	4 (57.1)	1.99 (0.60–6.53)	0.259			
6–9		20 (25.3)	20 (100.0)	136.86 (26.83–698.14) ^e	<0.001 ^e			

PFS, progression-free survival; HR, hazard ratio; S-GRAS, Stage, Grade, Resection status, Age, Symptom; CI, confidence interval.

^aS-GRAS components: European Network for the Study of Adrenal Tumour (ENSAT) stage (S), grading (G), resection status (R), age (A), and tumor- or hormone-related symptoms (S); ^bPatients subjected to S-GRAS analysis who underwent adrenalectomy and had available Ki-67 indices for S-GRAS scores; ^c*P* values comparing Harrell's C-index of grading with that of S-GRAS score; ^d*P* values comparing Harrell's C-index of grading with that of S-GRAS groups; ^eSignificant results (*P*<0.05).