

Supplemental Table S2. Baseline Characteristics of Total Thyroidectomy and Lobectomy Patients

Baseline variable	Total thyroidectomy (n=2,064)	Lobectomy (n=450)	P value
Demographic data			
Index year	2,012 (2,009–2,013)	2,014 (2,011–2,017)	<0.001
Age, yr	54 (50–60)	55 (52–60)	0.077
Sex			0.055
Women	1,710 (82.8)	355 (78.9)	
Men	354 (17.2)	95 (21.1)	
BMI, kg/m ²	24.1 (22.2–26.1)	23.75 (22.1–25.78)	0.109
Weight, kg	60 (54–67)	59 (55–67)	0.595
Income			0.564
Low	598 (29)	126 (28)	
Middle	492 (23.8)	118 (26.2)	
High	974 (47.2)	206 (45.8)	
Smoking			0.024
None	1,802 (87.3)	371 (82.4)	
Ex-smoker	139 (6.7)	41 (9.1)	
Current	123 (6)	38 (8.4)	
Alcohol consumption			0.031
None	1,529 (74.1)	327 (72.7)	
Mild to moderate	407 (19.7)	80 (17.8)	
Heavy	128 (6.2)	43 (9.6)	
Clinical data			
Systolic blood pressure, mm Hg	123 (113–132)	123 (112–132)	0.900
Diastolic blood pressure, mm Hg	78 (70–82)	78 (70–83)	0.322
Fasting blood sugar, mg/dL	95 (87–105)	95 (88–106.75)	0.320
Total cholesterol, mg/dL	199 (174–224)	199 (178–223)	0.697
Medical history			
Diabetes	511 (24.8)	108 (24)	0.781
Hypertension	895 (43.4)	188 (41.8)	0.574
Hyperlipidemia	940 (45.5)	241 (53.6)	0.002
Cerebrovascular disease	210 (10.2)	39 (8.7)	0.377
Cardiovascular disease	262 (12.7)	37 (8.2)	0.010

Values are expressed as median (interquartile range) or number (%).
BMI, body mass index.