

Supplementary Table S2. Correlation of TRG of KSP with clinicopathological parameters in rectal cancer.

	TRG of KSP		p-value
	TRG0, 1	TRG1, 2	
Age (yr)			
<65	16 (15.7)	34 (33.3)	.215
≥65	11 (10.8)	41 (40.2)	
Sex			
Male	18 (17.6)	52 (51.0)	.798
Female	9 (8.8)	23 (22.5)	
Differentiation			
Well	2 (2.0)	3 (3.0)	.635
Moderate	23 (23.2)	66 (66.7)	
Poorly	2 (2.0)	3 (3.0)	
Lymphovascular invasion			
Absent	19 (20.2)	52 (55.3)	.071
Present	2 (2.1)	21 (22.3)	
pT stage			
Tx–T2	24 (23.5)	26 (25.5)	<.001
T3–T4	3 (2.9)	49 (48.0)	
pN stage			
N0	24 (23.5)	47 (46.1)	.011
N1–N2	3 (2.9)	28 (27.5)	
pM stage			
M0	27 (26.7)	73 (72.3)	>.99
M1	0	1 (1.0)	
MSI			
MSS and MSI-L	24 (23.5)	69 (67.6)	.625
MSI-H	3 (2.9)	6 (5.9)	
Recurrence			
No	23 (22.8)	53 (52.5)	.225
Yes	4 (4.0)	21 (20.8)	
Death			
No	26 (25.5)	66 (64.7)	.509
Yes	1 (1.0)	9 (8.8)	
TRG of Mandard			
TRG1, 2	27 (26.5)	0	<.001
TRG3	0	37 (36.3)	
TRG4, 5	0	38 (37.3)	

TRG, tumor regression grade; KSP, The Korean Society for Pathologists; MSI, microsatellite instability; MSS, microsatellite stable; MSI-L, MSI-low; MSI-H, MSI-high.