

EXTREMELY WELL DIFFERENTIATED CARCINOMA

Supplementary Table S2. Immunohistochemical features of 19 cases of EWDA

Case No.	Histologic type	Immunohistochemistry						p53 ^b
		MUC5AC	MUC6	MUC2	CDX-2	c-Erb B2 ^a		
1	Gastric	+	Mucosa	-	-	-	-	1+
2	Gastric	+	Mucosa	-	-	2+	2+	
3	Gastric	+	Mucosa	-	-	-	2+	
4	Gastric	+	Mucosa	-	-	-	3+	
5	Gastric	+	Mucosa to submucosa	+, focal	-	-	1+	
6	Gastric	+	Mucosa to muscularis propria	-	-	-	3+	
7	Intestinal	Faint+	-	-	+	-	3+	
8	Gastric	+	Mucosa	-	-	-	3+	
9	Intestinal	Faint+	-	-	+	-	2+	
10	Gastric	+	Mucosa to submucosa	+, focal	+	-	2+	
11	Gastric	+	Mucosa to submucosa	+, focal	+; heterogeneous	-	3+	
12	Intestinal	Faint+	-	-	+	-	3+	
13	Gastric	+	Focal	-	Faint, focal +	2+	2+	
14	Gastric	+	-	+; focal	-	-	2+	
15	Intestinal	Faint+	-	-	+	-	2+	
16	Gastric	+	-	+; focal	-	-	2+	
17	Gastric	+	Mucosa	+; focal	-	-	2+	
18	Intestinal	Faint+	-	-	+	-	2+	
19	Intestinal	Faint+	-	-	+	-	2+	

EWDA, extremely well-differentiated adenocarcinoma.

^aC-Erb B2 grading followed the guidelines for HER2 testing issued by the College of American Pathologists; ^bp53 grading, 1+:10%~1/3; 2+, 1/3~2/3; 3+, >2/3.