

Supplementary Table 2. Studies on the translation of WCQ into Korean

Study	Subject	Subscales	NI	Type of populations (n/M or F)	Rating	RC		BT	CVT	
						OA	SS			
Han and Oh (1990)	Parent	Problem-focused/wishful thinking	8/5	Mothers of chronically ill children (120/F)	4 (1-4)	0.79	None	None	None	
		Detachment/seeking social support	6/7							
		Focusing on the positive/tension-reduction	4/3							
		Total items	33							
Park and Lee (1992)	Married women	Problem-solving/tension-reduction	15/9	Married women (303/F)	4 (0-3)	0.92	None	None	Done	
		Seeking social support	6							
		Avoidance/wishful thinking	4/5							
		Total items	39							
Kim and Lee (1985)	Colleges students	Problem-focused	17	College Students (125/M, 51; F, 74)	4 (0-3)	None	0.90	None	Done	
		Distancing/wishful thinking	10/5							0.85/0.72
		Seeking social support	5							0.74
		Distraction/bargaining	6/5							0.74/0.69
		Trying challenge/anger expression	5/3							0.45/0.50
Total items	54									
Kim (1987)	Colleges students	AC	21	College freshmen (803/M, 426; F, 377)	4 (0-3)	None	0.95	None	Done	
		Tension-reduction	23							0.94
		PC	12							0.90
		Seeking social support	6							0.87
		Total	62							
Son et al. (1999)	None	AC/PC	15/1		4 (0-3)	None	None	None	None	
		Total items	5							30

Abbreviations: AC, active coping; BT, back translation; CVT, construct validity test; F, female; M, male; NI, numbers of items; OA, overall; PC, passive coping; RC, reliability coefficient; SS, subscales; WCQ, ways of coping questionnaire.