

# Obstetrics & Gynecology Science

Kyu Hee Cho, et al. Single-port laparoscopic staging in ovarian cancer

**Supplementary Table 1.** Patient characteristics of single-port laparoscopic surgical staging (SPLS) and staging laparotomy group after propensity score matching (n=42)

Characteristics	SPLS (n=21)	Staging laparotomy (n=21)	P-value
Age (yr)	47.3 (20–85)	49.8 (19–84)	0.358 <sup>a)</sup>
BMI (kg/m <sup>2</sup> )	24.7 (18.7–36.8)	22.4 (16.7–28.5)	0.131 <sup>a)</sup>
Pelvic adhesion	12 (57.1)	8 (38.1)	0.217 <sup>b)</sup>
Tumor size (cm)	12.6 (6.2–21)	12.1 (6–18)	0.753 <sup>a)</sup>
FIGO stage			0.047 <sup>b)</sup>
IA	15 (71.4)	8 (38.1)	
IB	0 (0.0)	0 (0.0)	
IC1	1 (4.8)	4 (19.0)	
IC2	5 (23.8)	5 (23.8)	
IC3	0 (0)	4 (19.0)	
Histology			0.536 <sup>b)</sup>
Serous	3 (14.3)	5 (23.8)	
Mucinous	5 (23.8)	1 (4.8)	
Seromucinous	1 (4.8)	3 (14.3)	
Endometrioid	2 (9.5)	3 (14.3)	
Clear cell	6 (28.6)	6 (28.6)	
Malignant germ cell	1 (4.8)	2 (9.5)	
Sex cord-stromal	2 (9.5)	1 (4.8)	
Others	1 (4.8)	0 (0.0)	
Tumor grade			0.979 <sup>b)</sup>
Grade 1	5 (23.8)	4 (19.0)	
Grade 2	4 (19.0)	4 (19.0)	
Grade 3	6 (28.6)	6 (28.6)	
N/A	6 (28.6)	7 (33.3)	

Data are corrected using propensity score matching. Data are presented as the median (range) or number (%).

BMI, body mass index; N/A, not applicable.

<sup>a)</sup>Calculated using the Mann–Whitney nonparametric test; <sup>b)</sup>Calculated using the  $\chi^2$  parametric test.