

# Obstetrics & Gynecology Science

Yuki Yamada, et al. Tumor laterality in ovarian cancer

**Supplementary Table 1.** Clinicopathological characteristics of the patients according to site of origin

	<b>Right ovary (n=40)</b>	<b>Left ovary (n=33)</b>	<b>P-value</b>
Age (yr)	61.0 (38-86)	62.0 (40-90)	0.363
BMI (kg/m <sup>2</sup> )	22.5 (16.3-35.2)	21.8 (16.8-30.7)	0.855
FIGO stage (2014)			
III	29 (72.5)	21 (63.6)	
IV	11 (27.5)	12 (36.4)	0.417
Histologic subtype			
Non-serous	18 (45.0)	11 (33.3)	0.311
Serous	22 (55.0)	22 (66.7)	
CA125	963.5 (33-18,338)	358 (11-6,065)	0.085
Ascites formation			
No	12 (30.0)	9 (27.3)	0.798
Yes	28 (70.0)	24 (72.7)	
Residual tumor			
No	11 (27.5)	10 (30.3)	0.792
Yes	29 (72.5)	23 (69.7)	
Tumor diameter (mm)	90 (34-335)	118 (46-188)	0.012

Values are presented as median (range) or number of patients (%).

BMI, body mass index; FIGO, the International Federation of Gynecology and Obstetrics; CA125, cancer antigen 125.