

Supplementary Table 2. Univariate and multivariate analysis of prognostic factors in revised stage III-IV ovarian cancer

	Patients	Univariate analysis			Multivariate analysis		
		Hazard ratio	95% CI	P-value	Hazard ratio	95% CI	P-value
Age (yr)	<55	39	Ref				
	≥55	92	0.454	0.745-1.932	0.355		
BMI (kg/m^2)	<18.5	25	Ref				
	≥18.5	106	1.648	0.910-2.985	0.096		
Revised FIGO stage	III	50	Ref			Ref	
	IV	81	2.452	1.441-4.172	0.001	2.119	1.219-3.683
Histologic subtype	Non-serous	61	Ref				
	Serous	70	1.045	0.674-1.619	0.845		
CA125	<274	43	Ref				
	≥274	83	1.259	0.741-2.173	0.367		
Ascites formation	No	34	Ref				
	Yes	97	1.49	0.799-2.782	0.207		
Residual tumor	No	23	Ref			Ref	
	Yes	108	2.543	1.159-5.580	0.016	1.879	0.829-4.260
Tumor diameter (mm)	>50.5	22	Ref				
	≥50.5	109	1.845	0.797-4.271	0.383		

CI, confidence interval; BMI, body mass index; FIGO, the International Federation of Gynecology and Obstetrics, CA125, cancer antigen 125.