

**Supplementary Table 4.** Enhancement and Echogenic Texture on Contrast-Enhanced Harmonic Endoscopic Ultrasonography and on Conventional Endoscopic Ultrasonography for Gallbladder Mass

	Enhancement texture		Echogenic texture	
	Homogenous	Inhomogenous	Homogenous	Inhomogenous
Benign	61	49	91	19
Malignant	1	34	8	27
Total	67	83	99	46

**Supplementary Table 5.** Univariate and Multivariate Analysis of Variables for Gallbladder Neoplasms

Variables	Univariate		Multivariate			
	Odds ratio (95% CI)	p-value	Conventional EUS		CEH-EUS	
			Odds ratio (95% CI)	p-value	Odds ratio (95% CI)	p-value
Sex	1.142 (0.583–2.238)	0.699				
Age	1.079 (1.047–1.112)	<0.001	1.038 (1.002–1.074)	0.037	1.037 (1.002–1.074)	0.040
Hypoechoic foci	20.222 (7.506–54.485)	<0.001	4.993 (1.414–17.629)	0.012	5.212 (1.658–16.377)	0.005
Tumor size	1.100 (1.057–1.145)	<0.001	1.046 (0.992–1.103)	0.097	1.049 (0.997–1.105)	0.067
Multi-lobulated	2.991 (1.451–6.164)	0.003	1.893 (0.697–5.146)	0.211	1.788 (0.655–4.879)	0.256
Multiple lesions	10.537 (2.428–45.723)	0.002	4.039 (0.837–19.495)	0.082	4.362 (0.901–21.119)	0.067
ConvEUS inhomogeneous	7.071 (3.337–14.980)	<0.001	1.512 (0.461–4.952)	0.495		
CEH-EUS inhomogeneous	4.000 (1.831–8.737)	0.001			1.802 (0.689–4.717)	0.230

EUS, endoscopic ultrasonography; CEH-EUS, contrast-enhanced harmonic endoscopic ultrasonography; CI, confidence interval.