

Supplementary Table 3. Body composition of patients according to the neurologic outcome at discharge in subgroups based on sex

| Variable | Good neurologic outcome at discharge | Poor neurologic outcome at discharge | P-value |
|-------------------------------------|--------------------------------------|--------------------------------------|---------|
| Male | (n=23) | (n=21) | |
| Proportion of fat mass | 18.3 (15.6–22.2) | 19.6 (17.0–22.1) | 0.630 |
| Proportion of total skeletal muscle | 45.9 (43.1–47.3) | 44.6 (41.9–47.2) | 0.692 |
| Proportion of upper limb muscle | 8.4 (7.6–8.9) | 8.6 (7.6–10.2) | 0.401 |
| Proportion of lower limb muscle | 27.0 (24.9–29.3) | 25.4 (23.0–27.8) | 0.316 |
| Female | (n=3) | (n=19) | |
| Proportion of fat mass | 33.5 (33.0–35.0) | 30.0 (22.7–40.0) | 0.651 |
| Proportion of total skeletal muscle | 35.5 (34.3–35.7) | 37.3 (30.7–40.4) | 0.651 |
| Proportion of upper limb muscle | 5.0 (4.7–5.3) | 5.7 (5.0–8.0) | 0.308 |
| Proportion of lower limb muscle | 22.5 (21.3–25.0) | 20.8 (17.1–22.8) | 0.308 |

Values are presented as median (interquartile range).