

## Implant therapy in oral and maxillofacial fields

Young-Kyun Kim, D.D.S., Ph.D.

Editor-in-Chief of JKAOMS

*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Seoul National University Bundang Hospital, Seongnam, Korea*

Dental implants have been established as a universal field of dental care. Dental implants are considered first when giving treatment for edentulous areas with severe bone deficiency, as well as when a single tooth is missing. Dental implant therapy is accompanied by a combination of oral and maxillofacial surgery, periodontology, and prosthodontic treatment. It is necessary to plan for upper prosthetic treatment before implant surgery. A top-down treatment plan is a basic principle of implant therapy. The oral and maxillofacial surgeon should consult with the prosthodontist and obtain a consensus on the implant treatment before surgery<sup>1,2</sup>. Some oral and maxillofacial surgeons have almost no knowledge of prosthodontic treatment or may even neglect the treatment. Implant placement for which upper prosthetic treatment cannot be properly performed may be entirely ineffective. Some operators believe that severe bone defects make implant misplacement inevitable. Even if an implant is successfully placed during major reconstructive surgery, it may be ineffective if it permits no upper prosthetic treatment or leads to improper prosthesis. Another consideration commonly neglected by oral and maxillofacial surgeons is implant maintenance care. Implant treatment should not be started if there are no postsurgical concerns about prosthodontic treatment or periodontal maintenance care. Dental implants are not considered to be an area in a specific department of professional care, but instead are a field that requires an integrative idea

of dental care. We stress here that oral and maxillofacial surgeons also need to have positive concern about prosthodontic treatment, periodontal treatment, and about implant maintenance care so that they can prevent their own field of oral and maxillofacial surgery from becoming irrelevant with regards to dental implants.

### Conflict of Interest

No potential conflict of interest relevant to this article was reported.

### References

1. Garber DA, Belser UC. Restoration-driven implant placement with restoration-generated site development. *Compend Contin Educ Dent* 1995;16:796, 798-802, 804.
2. Kim YK, Hwang JW, Lee HJ, Yeo IS, Yun PY. Influence of implant misplacement on the success of the final prosthesis: subjective evaluation by a prosthodontist of dental implants placed by an oral and maxillofacial surgeon. *J Korean Assoc Oral Maxillofac Surg* 2009;35:437-41.

---

#### **Young-Kyun Kim**

*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Seoul National University Bundang Hospital, 82 Gumi-ro 173beon-gil, Bundang-gu, Seongnam 13620, Korea*

TEL: +82-31-787-7541 FAX: +82-31-787-4068

E-mail: [kyk0505@snubh.org](mailto:kyk0505@snubh.org)

ORCID: <http://orcid.org/0000-0002-7268-3870>

©This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2015 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.