

Supplementary Table 1. Comparison of pain location and symptoms other than pain

Variable	Overall (n = 79)	No-antithrombotic therapy (n = 30)	Antithrombotic therapy (n = 49)	P-value
Pain location	56	21	35	0.538 ^{a)}
Epigastric	23 (41.1)	11 (52.4)	12 (52.2)	
Perumbilical	13 (23.2)	3 (14.3)	10 (28.6)	
Back pain	1 (1.8)	1 (4.8)	0 (0)	
Left abdomen	5 (8.9)	1 (4.8)	4 (11.4)	
Right abdomen	3 (5.4)	1 (4.8)	2 (5.7)	
Lower abdomen	1 (1.8)	1 (4.8)	0 (0)	
Ill defined ^{c)}	10 (17.9)	3 (14.3)	7 (20)	
Other symptom ^{d)}	22 (27.8)	13 (44.8)	9 (18.4)	0.013 ^{b)}
Nausea	16 (20.2)	9 (31)	7 (14.9)	0.096 ^{a)}
Vomiting	7 (8.8)	4 (13.8)	3 (6.4)	0.417 ^{a)}
Bloody stool	2 (2.5)	0 (0)	2 (4.3)	0.522 ^{a)}
Diarrhea	6 (7.6)	4 (13.8)	2 (4.3)	0.194 ^{a)}
Dyspepsia	1 (1.3)	1 (3.4)	0 (0)	0.382 ^{a)}

Values are presented as number (%).

^{a)}Fisher exact test, ^{b)}chi-square test. ^{c)}When patient cannot exactly locate their location of pain. ^{d)}Symptoms other than pain.