

Supplementary Table 2. Comparison of clinical outcomes and morphologic changes of symptomatic SISMAD patients (n = 63) by antithrombotic use

Variable	No-antithrombotic therapy (n = 23)	Antithrombotic therapy (n = 40)	P-value
Clinical outcome (n = 63)			
Hospital stay (day)	5.1 ± 1.3 (2–7)	7.7 ± 4.3 (3–28)	0.002 ^{a)}
Follow-up period (mo)	10.6 ± 10.3 (4–34)	27.22 ± 25.3 (3–86)	0.021 ^{a)}
Imaging follow-up period ^{c)} (mo)	10.4 ± 10.4 (3–34)	29.4 ± 25.9 (3–86)	0.022 ^{a)}
Pain resolution time (hr)	61 ± 30.2 (1–120)	49.7 ± 25.4 (10–96)	0.622 ^{a)}
Morphologic changes on follow-up imaging			
Remodeling (complete or incomplete)	19 (82.6)	33 (82.5)	0.991 ^{b)}
Maximal remodeling time (mo)	5.5 ± 3 (3–13)	11.8 ± 14.7 (1–80)	0.147 ^{a)}

Values are presented as mean ± standard deviation (range) or number (%).

SISMAD, spontaneous isolated superior mesenteric artery dissection.

^{a)}Mann-Whitney test, ^{b)}chi-square test. ^{c)}Time interval between initial diagnosis and last imaging follow-up.