

Supplementary Table 3. Comparison of clinical characteristics and angiographic features among symptomatic patients between remodeling and no remodeling group

Variable	Remodeling (n = 52)	No remodeling (n = 11)	P-value
Feature (n = 63)			
Age (yr)	51.3 ± 6.9 (36–72)	59.9 ± 13.4 (44–78)	0.005 ^{a)}
Male sex	43 (82.7)	8 (72.7)	0.425 ^{b)}
Comorbidity			
Hypertension	11 (21.2)	1 (9.1)	0.674 ^{b)}
Dyslipidemia	7 (13.5)	0 (0)	0.337 ^{b)}
Diabetes mellitus	0 (0)	0 (0)	NA
Ischemic heart disease	1 (1.9)	0 (0)	0.825 ^{b)}
Cerebrovascular disease	3 (5.8)	0 (0)	0.557 ^{b)}
Current or ex-smoker	24 (46.2)	1 (9.1)	0.039 ^{b)}
History of cancer	2 (3.8)	0 (0)	0.679 ^{b)}
Clinical feature			
Pain onset to treatment (hr)	58.9 ± 89.7 (0.5–480)	102 ± 187.4 (2–480)	0.912 ^{a)}
Angiographic feature			
SMA type			<0.001 ^{b)}
1	1 (1.9)	5 (45.5)	
2a	0 (0)	1 (9.1)	
2b	45 (86.5)	5 (45.5)	
3	6 (11.5)	0 (0)	
SMA length			0.053 ^{b)}
A	9 (17.3)	4 (36.4)	
B	28 (53.8)	7 (63.6)	
C	15 (28.8)	0 (0)	
Previous antithrombotics	5 (9.6)	1 (9.1)	0.957 ^{b)}

Values are presented as number only, mean ± standard deviation (range), or number (%).

NA, not applicable; SMA, superior mesenteric artery.

^{a)}Mann-Whitney test, ^{b)}Fisher exact test, ^{c)}chi-square test.