Ann Surg Treat Res 2022;103(6):360-371
https://doi.org/10.4174/astr.2022.103.6.360

Supplementary Table 3. The questionnaires for patient's satisfaction with treatment (n = 42)

Satisfaction for treatments	Visit 2 (1st week)			Visit 3 (2nd week)			Visit 4 (4th week)		
	Anal care (n = 22)	Control (n = 20)	P-value	Anal care (n = 22)	Control (n = 20)	P-value	Anal care (n = 22)	Control (n = 20)	P-value
Q1. Perianal dermatitis improvement	78.2	71.0	0.090	80.0	68.0	0.003	81.8	73.0	0.025
Q2. Anal pain	80.0	69.0	0.005	80.0	67.0	<0.001	81.0	72.0	0.008
Q3. Anal pruritus	76.4	66.0	0.009	79.1	69.0	0.003	78.2	72.0	0.115
Q4. Discomfort in bowel movements	72.7	66.0	0.170	71.8	68.0	0.306	73.6	67.0	0.088
Q5. Discomfort in everyday life	72.7	63.0	0.014	73.6	64.0	0.008	75.5	69.0	0.087
Q6. Overall quality of life	74.6	62.0	0.002	72.7	65.0	0.022	74.6	71.0	0.237
Q7. Satisfaction of anal pain care	71.8	59.0	0.010	75.5	62.0	<0.001	70.0	68.0	0.423

Only anal care group	Visit 2 (1st week)	Visit 3 (2nd week)	Visit 4 (4th week)	P-value
Q8. Convenience of use of skin protectants	79.1	81.0	82.9	0.124
Q9. Continuation of current treatment	87.6	86.2	86.2	0.549

The result shows the mean of the Likert scale score converted to score of 100.