

Supplementary Table 1. Postoperative outcomes of paraaortic lymph node dissection

Variable	Total (n = 53)	M1a group (n = 27)	M1bc group (n = 26)	P-value
Surgical area of PALND				
Aortocaval	42 (79.2)	20 (74.1)	22 (84.6)	0.544
Right inferior vena cava	41 (77.3)	20 (74.1)	21 (80.8)	0.800
Left aorta	42 (79.2)	19 (70.4)	23 (88.5)	0.199
Right common iliac	21 (39.6)	9 (33.3)	12 (46.2)	0.501
Left common iliac	28 (52.8)	16 (59.3)	12 (46.2)	0.496
Surgical area of extra PALND				
Right renal	2 (3.7)	1 (3.7)	1 (3.8)	>0.999
Left renal	9 (16.9)	5 (18.5)	4 (15.4)	>0.999
Pelvic lateral side	10 (18.8)	7 (25.9)	3 (11.5)	0.324
Complication, CDC				0.662
II	1 (1.8)	1 (3.7)	0 (0)	
IIIa	2 (3.7)	1 (3.7)	1 (3.8)	
IIIb	3 (5.6)	2 (7.4)	1 (3.8)	
V	1 (1.8)	1 (3.7)	0 (0)	

M1a, patients with isolated paraaortic lymph node metastasis; M1bc, patients with metastasis of paraaortic lymph node and other distant organs; PALND, paraaortic lymph node dissection; CDC, Clavien-Dindo classification.