

Supplementary Table 2. Long-term outcome after treatment of ruptured AAA treatment

| Variable | EVAR group | OSR group | Crude HR (95% CI) | P-value | Adjusted HR (95% CI) | P-value |
|------------------------------|------------|-----------|-------------------|---------|----------------------|---------|
| All-cause mortality | | | | | | |
| No. of patients | 266 | 207 | | | | |
| PY | 1,537.8 | 1,196.7 | | | | |
| Mortality rate (n/100 PY) | 17.3 | 17.3 | 1.07 (0.89–1.29) | 0.458 | 1.17 (0.98–1.41) | 0.087 |
| Mortality at ≤3 mo | | | | | | |
| Event | 240 | 205 | | | | |
| PY | 100.3 | 69.2 | | | | |
| Mortality rate (n/100 PY) | 239.2 | 293.3 | 1.18 (0.98–1.42) | 0.082 | 1.29 (1.07–1.55) | 0.009 |
| Mortality at >3 mo | | | | | | |
| No. of patients | 354 | 235 | | | | |
| Event | 26 | 2 | | | | |
| PY | 1,437.5 | 1,127.5 | | | | |
| Mortality rate (n/100 PY) | 1.81 | 0.18 | 0.10 (0.02–0.42) | 0.002 | 0.12 (0.03–0.49) | 0.003 |
| Reintervention ^{a)} | | | | | | |
| Event | 31 | 7 | | | | |
| PY | 1,416.7 | 1,177.0 | | | | |
| Incidence (n/100 PY) | 2.19 | 0.59 | 0.29 (0.13–0.67) | 0.004 | 0.31 (0.14–0.70) | 0.005 |
| Dialysis ^{a)} | | | | | | |
| No. of patients | 484 | 312 | | | | |
| Event | 23 | 10 | | | | |
| PY | 1,401.9 | 1,077.3 | | | | |
| Incidence (n/100 PY) | 1.64 | 0.93 | 0.64 (0.31–1.35) | 0.244 | 0.67 (0.31–1.43) | 0.300 |

EVAR, endovascular aneurysm repair; OSR, open surgical repair; HR, hazard ratio; CI, confidence interval; PY, person-years.

^{a)}Considered death as a competing risk.