

Supplementary Table 1. Clinical characteristics of liver donors

Characteristic	Donor 1	Donor 2	Donor 3
Ethnicity	Asian	Asian	Asian
Sex	Male	Male	Male
Age (yr)	63	64	25
Body mass index (kg/m ²)	23.0	20.6	24.3
Diagnosis	Hilar cholangiocarcinoma	Liver metastasis (gastric cancer)	Hepatic adenoma
Operation	Hepaticojejunostomy and jejunojejunostomy Right hemihepatectomy	Laparoscopic S7 wedge resection of liver	Laparoscopic left lateral sectionectomy of liver (S2, 3)
Albumin (g/dL)	2.9	4.1	4.7
PT (INR)	1.04	1.00	0.94
Total bilirubin (mg/dL)	2.51	0.40	0.70
ALP (U/L)	226	84	117
AST (U/L)	48	27	19
ALT (U/L)	46	17	19
Hepatitis	None	Hepatitis B	None
Alcohol use	Yes	NA	No
Smoking	No	NA	Yes
Creatinine (mg/dL)	0.64	1.05	0.80
Specimen weight (g)	N/A	0.42	0.42
WBC (K/ μ L)	10,000	4,870	5,100
Hemoglobin (K/ μ L)	13.2	10.7	15.5
Platelet (K/ μ L)	198	185	291

INR, international normalized ratio; NA, not available.